arXiv:1301.6508v1 [math-ph] 28 Jan 2013

Levy-Loewner Evolution: Exact Results

Igor LOUTSENKO

Université Paris Diderot
e-mail: loutsenko@math.jussieu.fr

Oksana YERMOLAYEVA

Institut de Physique Théorique, Saclay
e-mail: oksana.yermolayeva@cea.fr

Abstract

Levy-Loewner evolution is a generalization of the Schramm-Loewner evolution where the branching is possible in course of growth process. We define a class of radial Levy-Loewner evolutions for which sets of points of the average means beta-spectrum can be found exactly. These are Loewner evolutions driven by Levy processes with N first fourier coefficients of probability distribution corresponding to those of the Brownian motion at a countable number of special values of temperature, while the rest of coefficients remains free.

1 Introduction

We start with a simple introduction to radial Levy-Loewner evolution (a good introduction to the chordal LLE can be found in [15], [16], for introduction to Levy processes see e.g. [1] and references therein).

Let us consider iterative conformal mappings $z = F_n(w)$ from the exterior of the unit disc in the w-plane to the exterior of a bounded, simply connected domain in the z-plane: The nth mapping is a composition of n elementary "spike" mappings $z = f_i(w, \delta t_i)$, $i = 1..n$

$$F_n(w) = F_{n-1}(f_n(w, \delta t_n)), \quad F_0(w) = w \tag{1}$$

where

$$f_n(w,t) = e^{\imath\varphi_n} h\left(e^{-\imath\varphi_n} w, t\right), \quad h(w,t) = e^t \frac{w + 1 + \sqrt{(w+1)^2 - 4e^{-t}w}}{2w} - 1 \tag{2}$$

The elementary mapping $z = f_n(w, \delta t_n)$ attaches a radial "spike" of the length $\sqrt{\delta t_n}\,(1 + O(\delta t_n))$ located at the angle φ_n to unit disc: Here, the point $w = e^{\imath\varphi_n}$ on the unit circle in the w-plane is mapped to the tip of the spike in the z-plane (see Figure 1).

The mapping $z = h(w,t)$ that attaches a spike to the disc at $\varphi = 0$ satisfies the simplest Loewner equation $\frac{\partial h(w,t)}{\partial t} = w \frac{\partial h(w,t)}{\partial w} \frac{w+1}{w-1}$ and as a consequence

$$\frac{\partial f_n(w,t)}{\partial t} = w \frac{\partial f_n(w,t)}{\partial w} \frac{w + e^{\imath\varphi_n}}{w - e^{\imath\varphi_n}}$$

This equation is invariant wrt any conformal transformation $f_n(w,t) \to F\left(f_n(w,t)\right)$ and therefore the iterative compound mapping F_n in (1) can be represented as a solution of the Loewner equation at time $t = \sum_{i=1}^{n} \delta t_i$:

$$F_n(w) = F\left(w, \sum_{i=1}^{n} \delta t_i\right), \quad F(w,0) = w,$$

$$\frac{\partial F(w,t)}{\partial t} = w \frac{\partial F(w,t)}{\partial w} \frac{w + e^{\imath L(t)}}{w - e^{\imath L(t)}} \tag{3}$$



Figure 1: The elementary mapping $z = f_n(w, \delta t_n)$ (left) and the compound iterative mapping $z = F_n(w)$ (right) from exterior of unit circle in w-plane to exterior of a simply connected domain in z-plane.

where $L(t)$ is the piecewise constant function

$$L(t) = \varphi_i, \quad \sum_{j=1}^{i-1} \delta t_j < t < \sum_{j=1}^{i} \delta t_j$$

One can consider continuous-time Loewner evolutions as limits of the above iterative processes when $n \to \infty$ and $\delta t_n \to 0$. This iterative picture will be useful for the simplest derivation of linear integro-differential equation for moments of derivatives of conformal mappings for the Levy-Loewner evolution by the method proposed by M. Hastings [8].

We are interested in the case when $L(t)$ is a stochastic process without drift. Without loss of generality we set

$$\langle L(t) \rangle = 0, \tag{4}$$

where $\langle\rangle$ denote expectation (ensemble average).

When stochastic process $L(t)$ is continuous in time, conformal mapping $z = F(w, t)$ describes growth of a random continuous curve $\Gamma = \Gamma(t)$ starting from a point on a unit circle $|z| = 1$ at $t = 0$. On the other hand, when $L(t)$ is discontinuous in time, the growth branches in the z-plane.

If one requires that the Loewner evolution (3) is a conformally invariant Markovian process, in the sense that the time evolution is consistent with composition of conformal maps, then the necessary condition for such an evolution is that $L(t)$ must have independent stationary increments, i.e. $L(t)$ is a Markovian process with the probability distribution of $L(t+\tau) - L(t)$ depending only on τ. Among such processes there is a class of Levy processes considered in the present paper.

The only continuous (modulo uniform drift) process of Levy type is the Brownian motion

$$L(t) = B(t)$$

The Brownian motion is characterized by a single parameter - "temperature" κ :

$$\langle (B(t+\tau) - B(t))^2 \rangle = \kappa |\tau|. \tag{5}$$

Stochastic Loewner evolution driven by Brownian motion is called Schramm-Loewner Evolution (SLE, or SLE$_\kappa$). Since it describes non-branching planar stochastic curves with conformally-invariant probability distribution, SLE is a useful tool for description of boundaries of critical clusters in two-dimensional equilibrium statistical mechanics. In this picture, different κ correspond to different classes of models of statistical mechanics (good introduction to SLE for physicists can be found e.g. in [3], [6] as well as mathematical reviews can be found e.g. in [9], [10]).

By present, SLE is quite well studied and many exact results have been obtained. In particular, SLE is the only non-trivial example where multi-fractal spectrum as well as related β-spectrum of the curve is described exactly.

In contrast to SLE, there has been no much progress in study of branching processes: The branching growth processes are of great interest in non-equilibrium statistical mechanics, where extensive work in study of different two-dimensional growth models has been performed. In spite of great amount of numeric simulations, very few analytic results have been obtained in this direction so far.

In this article we present new analytic results for branching processes: These are examples of Levy-Loewner evolutions (LLE) where sets of points of the β-spectrum can be found exactly: The β-spectrum quantitatively describes subsets of the domain boundary where certain scaling laws apply [1] to derivatives of conformal mappings $F'(w)$ along the boundary. The integral means $\beta(q)$-spectrum of the domain is defined through the qth moment of derivative at the unit circle (i.e. at $|w| \to 1$) as follows

$$\beta(q) = \lim_{\epsilon \to 0+} \frac{\log \int_0^{2\pi} \langle |F'\left(e^{\epsilon + \imath\varphi}\right)|^q \rangle d\varphi}{-\log \epsilon} \tag{6}$$

The moments satisfy a linear integral equation which can be solved exactly for certain types of Levy processes for special sets of q. In the present work we exploit properties of these solutions to obtain exact values of β-spectrum on these sets.

Although the main part of the paper is devoted to the “bounded” whole-plane LLE, which is an infinite-time limit of the radial LLE introduced above, we consider in Section 6 and Appendices another “unbounded” version of the whole-plane LLE. The latter is an inversion $F(w,t) \to 1/F(1/w,t)$ of the bounded version and was studied due to its relationship with the problem of Bieberbach coefficients of conformal mappings for such processes [4], [12]: Expectations of the squares of the Bieberbach coefficients as well as two non-trivial points of the β-spectrum have been conjectured for two special examples of unbounded whole-plane LLEs in [4]. This conjecture has been proved in [12] for the first example and partially for the second one. The first example represents a point in an infinite set of exact values introduced in the present paper. Although the second example is also included in this set, it turns out to be in some sense an “exceptional” point. So, we complete the proof of the above mentioned conjecture for this exception in Appendix 2.

2 Equation for Moments of Derivative

To find $\beta(q)$-spectrum (6) one needs to estimate moments of derivative $\langle |F'\left(w,t\right)|^q \rangle$.

For this purpose it is convenient to change the variable w to $e^{\imath L(t)} w$ and consider the mapping

$$\tilde{F}(w,t) = F\left(e^{\imath L(t)} w, t\right)$$

for which the point $w = 1$ is the pre-image of the growing tip of the curve. Obviously,

$$\beta(q) = \lim_{\epsilon \to 0} \frac{\log \int_0^{2\pi} \tilde{\rho}\left(e^{\epsilon + \imath\varphi}, e^{\epsilon - \imath\varphi}; t|q\right), d\varphi}{-\log \epsilon},$$

where

$$\tilde{\rho}(w, \bar{w}; t|q) = \langle \left| \tilde{F}'\left(w,t\right) \right|^q \rangle. \tag{7}$$

---

[1] For introduction to multi-fractal analysis see e.g. [7], [2] and references therein

It turns out that the moments $\tilde{\rho}(w,\bar{w};t|q)$ satisfy an integral linear equation. The simplest derivation of such equation uses M. Hasting's iterative approach [8], [12]: In iterative picture (1) we consider the mappings

$$\tilde{F}_n(w) = \tilde{F}_{n-1}\left(e^{\imath(\varphi_n-\varphi_{n-1})}h(w,\delta t_n)\right).$$

By the chain rule

$$\left|\tilde{F}_n'(w)\right|^q = \left|\tilde{F}_{n-1}'\left(e^{\imath(\varphi_n-\varphi_{n-1})}h(w,\delta t_n)\right)\right|^q \left|h'(w,\delta t_n)\right|^q.$$

Taking expectations of the both sides of the above equation, with the account of (7), we get

$$\tilde{\rho}(w,\bar{w},t_{n-1}+\delta t_n) = |h'(w,\delta t_n)|^q \int_0^{2\pi} d\varphi P(\varphi,\delta t_n)\tilde{\rho}\left(e^{\imath\varphi}h(w,\delta t_n), e^{-\imath\varphi}\bar{h}(\bar{w},\delta t_n), t_{n-1}\right),$$

where $P(\varphi,t)$ is the probability that $L(t)=\varphi$ under condition that $L(0)=0$.

Taking into account that for small δt

$$h(w,\delta t) = w + w\frac{w+1}{w-1}\delta t,$$

in the first order in δt we obtain

$$\tilde{\rho}(w,\bar{w},t+\delta t) = \left(1+\delta t\left(w\frac{w+1}{w-1}\partial_w + \bar{w}\frac{\bar{w}+1}{\bar{w}-1}\partial_{\bar{w}} - \right.\right.$$
$$\left.\left. -\frac{q}{(w-1)^2} - \frac{q}{(\bar{w}-1)^2} + q\right)\right)\int_0^{2\pi} P(\varphi,\delta t)\tilde{\rho}\left(e^{\imath\varphi}w, e^{-\imath\varphi}\bar{w}, t\right)d\varphi,$$

so, in the $\delta t\to 0$ limit we arrive at the equation for $\tilde{\rho}$

$$\partial_t\tilde{\rho} = \left(-\hat{\eta} + w\frac{w+1}{w-1}\partial_w + \bar{w}\frac{\bar{w}+1}{\bar{w}-1}\partial_{\bar{w}} - \frac{q}{(w-1)^2} - \frac{q}{(\bar{w}-1)^2} + q\right)\tilde{\rho}. \tag{8}$$

Here the operator $\hat{\eta}$ acts on functions of $w,\bar{w}$ as follows

$$\hat{\eta}[\rho](w,\bar{w}) = \lim_{t\to 0}\frac{1}{t}\int_0^{2\pi}\left(\rho(w,\bar{w}) - \rho\left(e^{\imath\varphi}w, e^{-\imath\varphi}\bar{w}\right)\right)P(\varphi,t)d\varphi.$$

We consider only processes for which the above limit exists. On the unit circle, these Levy processes are defined by their characteristic exponents η_m

$$e^{-t\eta_m} = \langle e^{\mathrm{i}mL(t)}\rangle = \int_0^{2\pi} e^{\mathrm{i}m\varphi}P(\varphi,t)d\varphi$$

(i.e. $e^{-t\eta_m}$ is the Fourier transform of the probability distribution of $L(t)$).

According to the above definitions, the operator $\hat{\eta}$ acts diagonally on the basis of two-dimensional Taylor/Laurent expansions $w^n\bar{w}^m$, $n,m\in\mathbb{Z}$

$$\hat{\eta}[w^n\bar{w}^m] = \eta_{n-m}w^n\bar{w}^m. \tag{9}$$

Since we consider processes without drift, the characteristic exponents are real and symmetric

$$\eta_m = \bar{\eta}_m, \quad \eta_m = \eta_{-m}, \quad \eta_0 = 0.$$

In the case of Brownian motion (5), $P(\varphi,t)$ is a fundamental solution of the heat equation on the circle and

$$\eta_m = \frac{\kappa m^2}{2}, \quad \hat{\eta} = \frac{\kappa}{2}\left(w\partial_w - \bar{w}\partial_{\bar{w}}\right)^2, \tag{10}$$

i.e. equation (8) becomes the second order linear PDE.

3 The Whole Plane LLE

Equation (8) simplifies in the case of the whole plane LLE which is a properly scaled infinite-time limit of the radial LLE.

The whole-plane LLE $z = \mathcal{F}(w,t)$ can be viewed as the infinite-time limit of the radial evolution

$$\mathcal{F}(w,t) = \lim_{T\to\infty} e^{-T} F(w, T+t) \tag{11}$$

which describes growth of stochastic curve out of the point of origin in the plane. Obviously, $\mathcal{F}(w,t)$ also satisfies the radial Levy-Loewner equation (3).

Consider now the Laurent expansion of $\tilde{\mathcal{F}}(w,t) = \mathcal{F}\left(e^{\imath L(t)}w, t\right)$ at $w = \infty$

$$\tilde{\mathcal{F}}(w,t) = e^{t+\imath L(t)}\left(w + \sum_{i=0}^{\infty}\Phi_i(t) w^{-i}\right)$$

According to the above equation and (7)

$$\tilde{\rho} = e^{qt}\langle\left|1 - \sum_{i=1}^{\infty} i\Phi_i w^{-i-1}\right|^q\rangle = e^{qt}\rho, \quad \rho = \sum_{i=-1,j=-1}^{\infty}\frac{\rho_{i,j}}{w^{i+1}\bar{w}^{j+1}}, \quad \rho_{-1,-1} = 1 \tag{12}$$

By existence of limit (11), the probability distribution of $e^{-t-\imath L(t)}\tilde{\mathcal{F}}(w,t) = w + \sum_{i=0}^{\infty}\Phi_i(t)w^{-i}$ is time-independent [2]. So, in the case of the whole plane LLE, the function $\rho = e^{-qt}\tilde{\rho} = \langle\left|1 - \sum_{i=1}^{\infty} i\Phi_i w^{-i-1}\right|^q\rangle$ does not depend on time and

$$\frac{\partial\tilde{\rho}}{\partial t} = q\tilde{\rho}.$$

Therefore, for the whole-plane LLE, equation (8) becomes

$$L\rho = q\rho \tag{13}$$

$$L = -\hat{\eta} + w\frac{w+1}{w-1}\partial_w + \bar{w}\frac{\bar{w}+1}{\bar{w}-1}\partial_{\bar{w}} - \frac{q}{(w-1)^2} - \frac{q}{(\bar{w}-1)^2} + q \tag{14}$$

To find moments of derivatives we have to look for a non-vanishing and analytic at $w = \infty$ solution of the above equation. It is easy to see that such a solution is (up to a constant factor) unique.

Indeed, by analyticity of conformal mapping at $w = \infty$, the function $\rho(w,\bar{w})$ is analytic at infinity and $\rho(\infty) = 1$. Then substituting the expansion (12) into (13) we get the nine-term "two-dimensional" recurrence relation for the expansion coefficients ρ_{ij}

$$\sum_{l=0}^{2}\sum_{m=0}^{2} C_{ij}^{lm}\rho_{i-l,j-m} = 0 \tag{15}$$

with the following boundary conditions (see (12))

$$\rho_{-1,-1} = 1, \quad \rho_{i,j<-1} = \rho_{j<-1,i} = 0$$

and the recurrent coefficients (note the symmetry $C_{ij}^{lm} = C_{ji}^{ml}$)

$$C_{i,j}^{0,0} = -\eta_{i-j} - i - j - 2, \quad C_{i,j}^{1,1} = -4\eta_{i-j}, \quad C_{i,j}^{2,2} = -\eta_{i-j} + i + j - 2 - 2q,$$

$$C_{i,j}^{0,1} = 2\left(\eta_{i-j+1} + i + 1\right), \quad C_{i,j}^{0,2} = -\eta_{i-j+2} + j - i - 2 - q, \quad C_{i,j}^{1,2} = 2\left(\eta_{i-j+1} - j + 1 + q\right).$$

Any element ρ_{kn} can be found in a consecutive manner starting from $(i,j) = (-1,-1)$ and going along the "row" up to $j = n$ and repeating this procedure for consecutive rows up to $i = k$ by expressing ρ_{ij} as a linear combination of 8 elements: $\rho_{i-l,j-m}, l \in \{0,1,2\}, m \in \{0,1,2\}, (l,m) \neq (0,0)$. This fixes $\rho(w,\bar{w})$ uniquely. Therefore, once an analytic and non-vanishing at $w = \infty$ solution of (13) is found, it will correspond to the moments of derivatives.

---

[2] For details on existence of this infinite-time limit see Section 4.3 of [12] or Section 1.2 of [2]

Neither solution of (13), nor the β-spectrum can be found explicitly for general Levy processes, i.e. for arbitrary η_i. The only explicit example of the $\beta(q)$-spectrum is found for the SLE case: The dimension spectrum (which is a Legendre transform of the β-spectrum) was first predicted from quantum gravity by B.Duplantier [5] and then $\beta(q)$ was partially [3] proved in work by D.Belyaev and S. Smirnov [2] by using general facts from the theory of parabolic equations and that of harmonic measure. Below we find explicitly points of the $\beta(q)$-spectrum for the LLE driven by Levy processes with N first characteristic exponents η_i coinciding with that of Brownian motion at special values of κ

$$\eta_i = \frac{\kappa i^2}{2}, \quad i \leq N, \quad N = 1, 2, 3, \dots \tag{16}$$

and arbitrary characteristic coefficients for $i > N$.

In more details: For the sets of (q, κ) enumerated by integers N, n.

$$q = \frac{Nn(2N-n+1)}{N^2-n^2+n}, \quad \kappa = 2\frac{N-2n}{N^2-n^2+n}, \quad 0 \leq n \leq N/2 \tag{17}$$

and Levy processes (16), the β-spectrum equals

$$\beta = \kappa\frac{n^2}{2} = \frac{(N-2n)n^2}{N^2-n^2+n} \tag{18}$$

For a given κ, the maximal possible number of the above explicit points of the β-spectrum equals to the number of integer solutions (N, n) of equation $\kappa = 2\frac{N-2n}{N^2-n^2+n}$ with distinct $q = \frac{Nn(2N-n+1)}{N^2-n^2+n}$.

We obtain (18) by studying expansion of solution of (13) at $w \to \infty$ when (q, κ) belong to (17). Our procedure relies on solution of (13) along the infinite number of curves in the parametric (κ, q)-plane

$$q = \frac{M(2-\kappa M)(\kappa M+\kappa+2)(\kappa M+\kappa+4)}{2(2\kappa M+\kappa+2)^2}, \quad M = 1, 2, 3, \dots \tag{19}$$

The set (17) is a subset of (19) with $M = N - n$ and $\kappa = 2(M-n)/(M^2+2Mn+n)$.

4 SLE: Exact Solutions

To get the points of the β-spectrum for the Levy processes (16) we have to consider the whole-plane SLE.

Remind that in the case of the Loewner evolution driven by the Brownian motion (i.e. in the case of SLE) operator $\hat{\eta}$ is differential (see eq. (10)), and (13) becomes the second order linear PDE with

$$L = -\frac{\kappa}{2}\left(w\partial_w - \bar{w}\partial_{\bar{w}}\right)^2 + w\frac{w+1}{w-1}\partial_w + \bar{w}\frac{\bar{w}+1}{\bar{w}-1}\partial_{\bar{w}} - \frac{q}{(w-1)^2} - \frac{q}{(\bar{w}-1)^2} + q$$

It is convenient to make the following change of dependent variable ρ that eliminates the most singular terms in equation (13)

$$\rho = \left((1-w^{-1})(1-\bar{w}^{-1})\right)^{\gamma}\Theta, \quad q = 2\gamma + \frac{1}{2}\kappa\gamma - \frac{1}{2}\kappa\gamma^2, \tag{20}$$

---

[3] The analysys in [2] turns out not to be complete and is valid only for $q \geq -(\kappa+4)^2(\kappa+8)/128$, (see Appendix of [13])

Then (13) becomes

$$\Bigg(-\frac{\kappa}{2}(w-1)(\bar{w}-1)\left(w\frac{\partial}{\partial w}-\bar{w}\frac{\partial}{\partial \bar{w}}\right)^2+(\kappa\gamma-1)(w-\bar{w})\left(w\frac{\partial}{\partial w}-\bar{w}\frac{\partial}{\partial \bar{w}}\right)$$
$$+(w\bar{w}-1)\left(w\frac{\partial}{\partial w}+\bar{w}\frac{\partial}{\partial \bar{w}}\right)+\gamma V\Bigg)\Theta=0. \tag{21}$$

where

$$V=(w+\bar{w})\frac{\kappa+2}{2}+\kappa(\gamma-1)-2$$

Again, we can substitute the series expansion at $w=\infty$ for Θ

$$\Theta=\sum_{i=-1}^{\infty}\sum_{j=-1}^{\infty}\frac{\theta_{i,j}(\gamma,\kappa)}{w^{i+1}\bar{w}^{j+1}},\quad \theta_{-1,-1}=1 \tag{22}$$

into (21). This gives us the four term recurrence relation for expansion coefficients θ_{ij}

$$\sum_{k=0}^{1}\sum_{l=0}^{1}\mathcal{C}_{i,j}^{l,k}\theta_{i-l,j-k}=0,\quad \theta_{-1,-1}=1,\quad \theta_{i<-1,j}=\theta_{i,j<-1}=0, \tag{23}$$

$$\mathcal{C}_{i,j}^{0,0}=-\eta_{i-j}-i-j-2,\quad \mathcal{C}_{i,j}^{1,1}=-\eta_{i-j}+i+j+\gamma(\kappa\gamma-\kappa-2),$$
$$\mathcal{C}_{i,j}^{0,1}=\eta_{i-j+1}+(1-\kappa\gamma)(i-j+1)+\gamma\frac{\kappa+2}{2},\quad \mathcal{C}_{i,j}^{1,0}=\mathcal{C}_{j,i}^{0,1},$$
$$\eta_i=\frac{\kappa}{2}i^2$$

Note, that any matrix element $\theta_{i,j}$ can be found in a consecutive manner (starting from $\theta_{-1,-1}=1$) by expressing $\theta_{i,j}$ as a linear combination of 3 elements: $\theta_{i,j-1}$, $\theta_{i-1,j}$, $\theta_{i-1,j-1}$.

Moreover, since now the recurrence has four terms, the expansion coefficients θ_{ij} vanish for $|i-j|>M$, where $M=0,1,2\dots$, if the recurrence coefficient $\mathcal{C}_{i,j}^{0,1}$ vanishes at $j=i+M+1$. As follows from (23) and (20), such a $2M+1$-band truncation happens when

$$\kappa=2\frac{M-\gamma}{M^2+2M\gamma+\gamma},\quad q=\frac{\gamma(M+\gamma)(2M+1+\gamma)}{M^2+2M\gamma+\gamma} \tag{24}$$

or equivalently, when (19) holds.

The above equations determine infinite number of curves in the parametric (q,κ)-plane. The solutions of (21) along these curves have the form of the Fourier polynomials

$$\Theta(w,\bar{w})=\sum_{i=-M}^{M}w^{-i}f_i\left((w\bar{w})^{-1}\right), \tag{25}$$

where coefficients $f_i(\xi),\xi=1/(w\bar{w})$ are regular at $\xi=0$. Since $\theta(w,\bar{w})$ is real and symmetric

$$f_{-i}(\xi)=\xi^i f_i(\xi),\quad \bar{f}_i=f_i. \tag{26}$$

Note, that the "two-dimensional" recurrence relation (22) can be also rewritten in the form of "one-dimensional" three-term differential-difference equation by substituting the above Fourier expansion (25) into equation (21)

$$\xi A_{i+1}f_{i+1}+A_{-i+1}f_{i-1}+\left(B_i+(1-\xi)C_i\right)f_i+2\xi(\xi-1)\frac{df_i}{d\xi}=0, \tag{27}$$

where

$$A_i = \frac{\kappa}{2}(i-\gamma)^2 + i + \gamma + \frac{\kappa}{2}\gamma(1-\gamma), \quad B_i = -\kappa(i^2-\gamma^2+\gamma) - 2\gamma, \quad C_i = \frac{\kappa}{2}i^2 - \kappa\gamma^2 i + \kappa\gamma + 2\gamma$$

In this representation, the truncation condition is $A_M = 0$. When this truncation takes place, (27) becomes (with an account of the symmetry relation (26)) a system of the $M+1$ first order linear ODEs for $f_0(\xi), \dots, f_M(\xi)$. This system can be reduced to a linear ODE of the order $M+1$ with polynomial coefficients. [4] By asymptotic analysis of these ODEs one can find the β-spectrum along the curves (24) (or equivalently along (19)) [13].

We do not need to perform this analysis in this section, since the β-spectrum of this version of the whole-plane SLE can be derived for positive values of q using analysis of singularities of PDE (13) at $w = 1$ and general properties of the harmonic measure [2]. However, for an unbounded version of the whole-plane SLE which will be considered later in this work, the only found so far way of obtaining the β-spectrum for the values of q of interest is by direct solution or asymptotic analysis of the three-term relations [13].

5 LLE and SLE: Common Solutions

Now we are ready to obtain the main result (16-18). For this we consider solutions of (13) that are Fourier polynomials.

It follows from (25) and (20) that, when γ is integer and conditions (24) hold, ρ becomes the Fourier polynomial

$$\gamma = n, \quad \rho = \left(1 + \xi^{-1} - w^{-1} - w\xi^{-1}\right)^n \sum_{i=-M}^{M} w^i f_i(\xi) = \sum_{i=-N}^{N} w^i g_i(\xi), \quad n \in 0, 1, 2, \dots \tag{28}$$

and we have truncation of expansion (12) at $|i-j| = N$, i.e.

$$\rho_{i,j} = 0, \quad |i-j| > N. \tag{29}$$

Remind, that we consider LLE driven by Levy processes (16) with first N characteristic coefficients coinciding with that of Brownian motion at special values of κ (given by (17)). From the relation (15) for ρ_{ij} it follows that this truncation takes place independently of the values of the rest of characteristic coefficients $\eta_i, i > N$.

Indeed, taking the expansion coefficients for the SLE case (29), and substituting them into recurrence relation (15) it is easy to see that all recurrent coefficients containing η_l with $|l| > N$ are multiplied by zeros, since (29) holds. Thus only remaining terms in the recurrence equations coincide with those for the corresponding SLE case, and the truncated SLE solution is also solution for the corresponding LLE (16,17).

Therefore, when parameters (q, κ) take values (17), the β-spectrum of LLE (16) takes the values coinciding with those of the corresponding SLE-process. The β-spectrum of the whole-plane SLE is known and equals [2], [13]

$$\beta = \begin{cases} \kappa\frac{\gamma^2}{2} - 2\gamma - 1, & q \le -1 - \frac{3\kappa}{8} \\ \kappa\frac{\gamma^2}{2}, & -1 - \frac{3\kappa}{8} \le q \le \frac{3(\kappa+4)^2}{32\kappa} \\ q - \frac{(\kappa+4)^2}{16\kappa}, & q \ge \frac{3(\kappa+4)^2}{32\kappa} \end{cases}$$

where in terms of q and κ (see (20))

$$\gamma = \frac{\kappa + 4 - \sqrt{(\kappa+4)^2 - 8q\kappa}}{2\kappa}$$

---

[4] For instance, in the $M = 1$ case, (27) is reduced to the hypergeometric equation and $\Theta(w, \bar{w})$ is explicitly expressed in terms of hypergeometric functions when, according to (19), $q = 2(4-\kappa^2)(\kappa+1)/(3\kappa+2)^2$

Note, that q from the set (17) are always bounded within the range $-1-\frac{3\kappa}{8} \leq q \leq \frac{3(\kappa+4)^2}{32\kappa}$, and, therefore, values of the spectrum for this set equal $\kappa\gamma^2/2$. Since, according to (28), for such truncations $\gamma = n$, we arrive to the main result (16-18).

6 Unbounded Whole-Plane LLE

Remind that in the version of LLE considered above, the curve grows from the origin towards infinity. This version is called the bounded whole-plane LLE. There exist another, unbounded version of the whole-plane LLE, where (an unbounded) curve grows from the infinity towards the origin. Both versions are related by the inversion $\mathcal{F}(w,t) \to 1/\mathcal{F}(1/w,t)$. Obviously, the unbounded mapping satisfies the same Loewner equation (3), but now $\mathcal{F}$ maps interior of the unit disc into the complement of the unbounded curve in the plane

$$\mathcal{F}(w,t) = e^{-t}\left(w + \sum_{i=2}^{\infty} \mathcal{F}_i(t) w^i\right), \quad |w| < 1$$

So, we refer to the unbounded whole-plane LLE as the "interior problem", while the bounded version considered in the previous sections is called the "exterior problem".

By analogy with the exterior problem, the moments of derivatives

$$\tilde{\rho} = \langle |\mathcal{F}'(e^{iL(t)}w, t)|^q \rangle$$

depend on time only through exponential scaling, i.e. (note the difference in the exponent sign in comparison with (12))

$$\tilde{\rho} = e^{-qt}\rho$$

where

$$\rho(w,\bar{w};q) = \sum_{i=1,j=1}^{\infty} \rho_{ij} w^{i-1}\bar{w}^{j-1}, \quad \rho_{1,1} = 1 \tag{30}$$

is time-independent.

Similarly to the exterior problem, $\rho(w,\bar{w};q)$ satisfies an integral equation which now writes as follows

$$L\rho = -q\rho, \tag{31}$$

where L is given by (14). Now an analytic and non-vanishing at $w = 0$ solution of (31) corresponds to the moments of derivatives of the interior problem.

The $\beta(q)$-spectrum of interior problem equals

$$\beta(q) = \lim_{\epsilon \to 0+} \frac{\log \int_0^{2\pi} \rho(e^{-\epsilon + \imath\varphi}, e^{-\epsilon - \imath\varphi}; q) d\varphi}{-\log \epsilon} \tag{32}$$

It can be computed explicitly for interior whole-plane SLE [13], [14] along the countable number of curves in the parametric (q,κ)-plane, which include all points of the interest. It equals (see Appendix 1)

$$\beta = \begin{cases} \kappa\frac{\gamma^2}{2} - 2\gamma - 1, & q \leq -1 - \frac{3\kappa}{8} \\ \kappa\frac{\gamma^2}{2}, & -1 - \frac{3\kappa}{8} \leq q \leq Q(\kappa) \\ 3q - \frac{1}{2} - \frac{1}{2}\sqrt{1+2q\kappa}, & q \geq Q(\kappa) \end{cases} \tag{33}$$

where

$$Q(\kappa) = \frac{\kappa^2 + 8\kappa + 12 - 2\sqrt{2\kappa^2 + 16\kappa + 36}}{16\kappa} \tag{34}$$

In the interior problem, β-spectrum of SLE shows two transitions, one of which (similarly to the exterior case) happens at negative $q = -1 - \frac{3\kappa}{8}$ due to singularity (zero of derivative) of the conformal mappings at the tip of the curve. Another one happens at positive $q = Q(\kappa)$ due to singularity of the mapping at $z = \infty$.

By analogy with the previous sections, one can also explicitly find points of the β-spectrum for the LLE driven by Levy processes (16) with N first characteristic coefficients η_i coinciding with that of Brownian motion at special values of κ and arbitrary η_i for $i > N$.

In more details: The values of the β spectrum of interior problem can be found for processes (16) with

$$\kappa = 2\frac{N+2n}{N^2+n^2-n}, \quad q = \frac{nN(2N-n+1)}{N^2+n^2-n}, \quad n \leq N, \quad N, n = 0, 1, 2, \dots \tag{35}$$

These values equal

$$\beta = \frac{N(6Nn - 3n^2 - N + n)}{N^2+n^2-n} \tag{36}$$

Note that q and κ in (35) are such that q is always bigger than $Q(\kappa)$ in (34), i.e. (36) is obtained from the part of the SLE β-spectrum above the transition point at $q = Q(\kappa)$.

The $\beta(q)$-spectrum of SLE below the transition at $q = Q(\kappa)$ can be obtained by the method of [2]. Here, however we need spectrum for $q \geq Q(\kappa)$ which is so far determined with the help of exact solutions of (31) [14]. We outline the derivation of the spectrum for the interior whole-plane SLE in Appendix 1.

It is now convenient to make the following change of the dependent variable in equation (31)

$$\rho = ((1-w)(1-\bar{w}))^{\gamma}\,\Theta, \quad \Theta = \sum_{i=1,j=1} \theta_{ij} w^{i-1} w^{j-1}, \quad \theta_{1,1} = 1 \tag{37}$$

After this change we get the PDE of the form (21) with

$$V = (\kappa - \kappa\gamma + 6)w\bar{w} + \frac{2\kappa\gamma - \kappa - 6}{2}(w + \bar{w}) \tag{38}$$

Again, we are looking for solutions Θ that are Fourier polynomials

$$\Theta(w, \bar{w}) = \sum_{i=-M}^{M} w^i f_i(\xi), \quad \xi = w\bar{w} \tag{39}$$

Substituting (39) into (21), (38), we get the three-term recurrent relation for $f_i(\xi)$ of the form (27) with

$$A_i = \frac{\kappa}{2}(i-\gamma)^2 + i - 3\gamma - \frac{\kappa}{2}\gamma(1-\gamma), \quad B_i = -\kappa(i^2+\gamma^2-\gamma) + 6\gamma, \quad C_i = \kappa\frac{i^2 - 2\gamma + 2\gamma^2}{2} - i - 6\gamma. \tag{40}$$

Remind, that since $\Theta(w, \bar{w})$ is real and symmetric, the condition (26) holds. The $2M+1$-diagonal truncation (39) of solution Θ happens when $A_M = 0$. This condition defines the infinite countable set of curves in the parametric (q, κ)-plane

$$\kappa = 2\frac{M + 3\gamma}{M^2 + 2M\gamma + 2\gamma^2 - \gamma}, \quad q = \frac{\gamma(M+\gamma)(2M+1+\gamma)}{M^2 + 2M\gamma + 2\gamma^2 - \gamma} \tag{41}$$

In the case when γ is an integer, ρ is also the Fourier polynomial

$$\gamma = n, \quad \rho = \left(1 + \xi - w - \xi w^{-1}\right)^n \sum_{i=-M}^{M} w^i f_i(\xi) = \sum_{i=-N}^{N} w^i g_i(\xi)$$

Similarly to the exterior case, the set of parameters (35) corresponds to points on curves (41) where $\gamma = n$, and therefore $N = M + n$. Repeating arguments used for the exterior problem, one concludes that at these points the solution $\rho(w, \bar{w})$ for SLE coincides with that for LLE driven by Levy processes (16), (35) and so we prove the main result in the case of interior problem.

Note, that in the case of the interior problem one more explicit example exists: For Levy processes with $\eta_1 = 1$ and arbitrary $\eta_i, i > 1$ we have [5]

$$\beta(2) = 4, \quad \eta_1 = 1 \tag{42}$$

This point of the spectrum has been conjectured using computer algebra in [4]. It has been also conjectured that for the above values of parameters $\rho_{i,i} = i^3$. We prove this conjecture in Appendix 2.

7 Conclusion

In summary: From elementary consideration it follows that any analytic and non-vanishing at $w = \infty/w = 0$ solution of the equation for moments (13)/(31) is unique (up to a constant factor) and, therefore, provides a point in the β-spectrum of the exterior/interior whole plane LLE, respectively.

For the class of LLE driven by Levy processes with N first characteristic exponents equal to those of the Brownian motion and with the rest of characteristic exponents being free parameters, these solutions coincide with solutions of PDE for corresponding Schramm-Loewner Evolutions when (κ, q) belong to the countable sets of parameters (17) for the exterior problem or to the set (35) for the interior problem respectively. As a consequence, the corresponding β-spectra can be found exactly for the parameters of interest either by method by D.Belyaev and S.Smirnov in the case of exterior problem, or through analysis of exact solutions in the both exterior and interior problems.

For a given κ and N the number of explicit points (18) in the β spectrum equals the number of integer solutions (N', n) with distinct $q = \frac{N'n(2N'-n+1)}{N'^2-n^2+n}$ of the Diophantine equation

$$\kappa = 2\frac{N' - 2n}{N'^2 - n^2 + n}, \quad 0 \leq N' \leq N, \quad n \leq N'/2$$

in the case the exterior problem, and number of integer solutions of equation

$$\kappa = 2\frac{N' + 2n}{N'^2 + n^2 - n}, \quad 0 \leq N' \leq N, \quad n \leq N'$$

with distinct $q = \frac{nN'(2N'-n+1)}{N'^2+n^2-n}$ in the case of the interior problem (where β is given by (36)) respectively.

Acknowledgement

We would like to acknowledge help received from A. Boutet de Monvel, V. Spiridonov and A. Zhedanov. This work has been supported by the European Commission 7th framework IEF grants.

Appendix 1: β-spectrum of the interior whole-plane SLE

Let us briefly outline the derivation of (33), (34) on the infinite set of curves (41) (for more details see [13] or [14]).

The derivation is based on the asymptotic analysis [6] of the difference-differential equation (27), (40) for $|w| \to 1$ (i.e. at $\xi \to 1$): The system of $2M + 1$ linear ODEs (27), (40) for $f := \{f_i(\xi), i = -M..M\}$ has $2M + 1$ linearly independent solutions that behave at $\xi \to 1$ as

$$f(\xi) \to (1 - \xi)^{-\beta_l} \mathcal{V}_l,$$

---

[5] The sub-case of (42) with $\eta_1 = 1$, $\eta_2 = 4$ and arbitrary $\eta_i, i > 2$ is contained in the set (35-36) at $N = 2, n = 1$.

[6] There also exist explicit algebraic and hypergeometric solutions for the $M = 0$ and $M = 1$ curves (41) respectively (see [14])

where $\beta_l, l = 0..2M$, are $2M+1$ eigenvalues of a $2M+1 \times 2M+1$ tri-diagonal matrix and $\mathcal{V}_l$ is a ξ-independent $2M+1$ component eigenvector corresponding to the eigenvalue β_l. Indeed, expanding solutions of (27) at $\xi \to 1$ as follows

$$f_i(\xi) = (1-\xi)^{-\tilde{\beta}}\psi_i + \ldots, \quad i = -M..M \tag{43}$$

and substituting the above expansion into (27) we get the following three-term recurrence relation for ψ_i

$$R[\psi] = \tilde{\beta}\psi, \quad R[\psi]_i := \frac{1}{2}\left(A_{i+1}\psi_{i+1} + A_{-i+1}\psi_{i-1} + B_i\psi_i\right) \tag{44}$$

with $\tilde{\beta}$ being an eigenvalue of the $2M+1 \times 2M+1$ three-diagonal matrix R.

As follows from (39), (43), the Fourier transform $\Psi(\varphi)$ of ψ_j

$$\Psi(\varphi) := \sum_{j=-M}^{M} \psi_j e^{\imath j\varphi}, \quad \bar{\Psi} = \Psi \tag{45}$$

is the "angular part" of $\Theta(w, \bar{w})$ at the unit circle $w = e^{\imath\varphi}$, i.e.

$$\Psi(\varphi) = \lim_{r\to 1-} \frac{\Theta\left(re^{\imath\varphi}, re^{-\imath\varphi}\right)}{(1-r^2)^{\tilde{\beta}}} \tag{46}$$

According to definition of β-spectrum (32), as well as due to (37) and (46)

$$\beta = \tilde{\beta} + \lim_{\epsilon\to 0+} \frac{\log\int_0^{2\pi}\left((1-e^{-\epsilon+\imath\varphi})(1-e^{-\epsilon-\imath\varphi})\right)^{\gamma}\Psi(\varphi)d\varphi}{-\log\epsilon},$$

Then, from the fact that the factor $\left((1-e^{-\epsilon+\imath\varphi})(1-e^{-\epsilon-\imath\varphi})\right)^{\gamma}$ in (20) ceases to be integrable on the unit circle $w = e^{\imath\varphi}$ when $\gamma < -1/2$, from the previous equation and eq. (46) we have

$$\beta = \left\{ \begin{array}{ll} \tilde{\beta} - 2\gamma - 1, & \gamma \le -\frac{1}{2} \\ \tilde{\beta}, & \gamma \ge -\frac{1}{2} \end{array} \right., \quad \tilde{\beta} \in \{\beta_l, l = 0..2M\}, \tag{47}$$

provided $\Psi(\varphi)$ does not have zeros at $\varphi = 0$ when $\gamma < -1/2$ (which is the case). Therefore, the β-spectrum of the whole-plane SLE is determined by an eigenvalue of the three diagonal matrix (44).

The eigenvalues $\{\beta_l, l = 0..2M\}$ of the three-diagonal $2M+1 \times 2M+1$ matrix (44), (40) can be found either directly [11] or by recasting three-term recurrence relation (44), (40) in the form of the second order linear ODE for $\Psi(\varphi)$ which is amenable to hypergeometric form [13], [14]. The only even solutions of this equation are

$$\Psi = (1-\cos(\varphi))^{l/2}{}_2F_1\left(\frac{l}{2} - M, \frac{l}{2} + \gamma\frac{3M+1+4\gamma}{M+3\gamma}; \frac{1}{2}\middle|\frac{1+\cos(\varphi)}{2}\right), \quad l = 0, 2, 4, \ldots 2M \tag{48}$$

corresponding to even eigenvalues

$$\beta_l = \frac{2(M+3\gamma)\gamma^2 - (2M^2 + M - 8\gamma^2 + \gamma)l + (M+3\gamma)l^2}{2(M^2 + 2M\gamma + 2\gamma^2 - \gamma)}, \quad l = 0, 2, ..2M$$

The biggest eigenvalue, depending on M and γ, is either

$$\beta_0 = \frac{M+3\gamma}{M^2 + 2M\gamma + 2\gamma^2 - \gamma}\gamma^2 = \frac{\kappa\gamma^2}{2} \tag{49}$$

or

$$\beta_{2M} = \frac{(M+\gamma)(6M\gamma - M + 3\gamma^2)}{M^2 + 2M\gamma + 2\gamma^2 - \gamma} = 3q - \frac{1}{2} - \frac{1}{2}\sqrt{1 + 2q\kappa} \tag{50}$$

where (41) has been taken into account to express M and γ in terms of q and κ.

By non negativity of the moments of derivative, $\Psi(\varphi)$ must be non-negative on the unit circle $0 \leq \varphi \leq 2\pi$. For $q > Q(\kappa)$ the only non-negative $\Psi(\varphi)$ in (48) corresponds to $l = 2M$ (see [13] or [14]). For $q < Q(k)$ the $\beta = \beta_0$, which can be shown by method by D. Belyaev and S.Smirnov [2]. Thus, the value of $\tilde{\beta}$ equals the maximal eigenvalue. [7] As a consequence one can observe transition in the means β-spectrum at points of the parametric (q, κ)-plane where $\beta_0 = \beta_{2M}$. These two eigenvalues are equal at the transition point $q = Q(\kappa)$ (see Eqs (33-34)).

Taking (47) into account, from (49), (50), one obtains $\beta(q)$-spectrum (33) along a countable number of curves (41) in the parametric (q, κ) plane for the interior whole-plane SLE.

Appendix 2: Interior problem: Proof for $\eta_1 = 1, q = 2$ case

As mentioned in the Introduction and Section 6, this Appendix completes proof of the conjecture by B. Duplantier et al [4], so we formulate its result in the form of the following

Theorem: For the interior whole plane LLE driven by Levy processes with $\eta_1 = 1$, for $q = 2$:

$$\rho_{ii} = i^3, \quad \beta(2) = 4$$

Proof:

In difference with the main examples considered above, the case $\eta_1 = 1$, $q = 2$ with arbitrary $\eta_i, i > 1$ does not, in general, show a finite-diagonal truncation of $\rho_{i,j}$. Representing $\rho(w, \bar{w})$ in the form

$$\rho(w, \bar{w}) = (1 - w)(1 - \bar{w})\Theta(w, \bar{w}) \tag{51}$$

from (31) with $q = 2$ we get

$$-\hat{\eta}[(1 - w)(1 - \bar{w})\Theta] + (w + 1)(\bar{w} - 1)w\frac{\partial\Theta}{\partial w} + (\bar{w} + 1)(w - 1)\bar{w}\frac{\partial\Theta}{\partial\bar{w}} + 3(2w\bar{w} - w - \bar{w})\Theta = 0 \tag{52}$$

where Θ is the series

$$\Theta = \theta_0(\xi) + \sum_{i=1}^{\infty}\left(w^i + \xi^i w^{-i}\right)\theta_i(\xi), \quad \xi = w\bar{w} \tag{53}$$

Substituting it into (52) we will get a three-term differential recurrence relation for $\theta_i(\xi)$ which does not truncate in general [8]. However, it turns out that in this recurrence relation, two equations defining functions $\theta_0(\xi)$ and $\theta_1(\xi)$ do not involve others $\theta_i(x), i \neq 0, 1$ when $\eta_1 = 1$.

Indeed, making the change of variable $\bar{w} = \xi/w$, then taking into account the facts that, according to (9), operator $\hat{\eta}$ commutes with $\xi = w\bar{w}$ and that $\hat{\eta}[w^i] = \eta_i w^i$, after substitution of series (53) into (52) we will get a series in w with coefficients depending on ξ. Coefficients of w^i define the following three-term differential recurrence relations

$$2\xi(\xi-1)\theta_i'(\xi)-(\eta_i + i + (\eta_i - i - 6)\xi)\,\theta_i(\xi)+\xi(\eta_i+i-2)\theta_{i+1}(\xi)+(\eta_i-i-2)\theta_{i-1}(\xi) = 0, \quad \theta_{-i}(\xi) = \xi^i\theta_i(\xi)$$

For $i = 0$ and $i = 1$ they write

$$(\xi-1)\theta_0'(\xi)+3\theta_0(\xi)-2\theta_1(\xi) = 0, \quad (1-\eta_1)\xi\theta_2(\xi)+2\xi(\xi-1)\theta_1'(\xi)+(7\xi-\eta_1\xi-\eta_1-1)\theta_1(\xi)+(\eta_1-3)\theta_0(\xi) = 0$$

When $\eta_1 = 1$, these equations do not involve θ_2:

$$(\xi - 1)\theta_0'(\xi) + 3\theta_0(\xi) - 2\theta_1(\xi) = 0, \quad \xi(\xi - 1)\theta_1'(\xi) + (3\xi - 1)\theta_1(\xi) - \theta_0(\xi) = 0$$

---

[7] The same conclusions can be also obtained for the values of interest from general properties of β-spectra [14]

[8] Truncation happens when $\eta_i = i + 2$

Since function ρ is analytic at $w = 0$, as well as $\rho(w = 0) = 1$, we have to look for a solution of the above system of two ODEs, such that $\theta_0(0) = 1$ and $\theta_1(\xi)$ is finite at $\xi = 0$. With these conditions we get

$$\theta_0 = \frac{1+\xi}{(1-\xi)^4}, \quad \theta_1 = -\frac{1}{(1-\xi)^4}$$

Taking into account (30), (53) and (51) from the above we obtain that $\rho_{ii} = i^3$, if $\eta_1 = 1$. Then from (32) it follows that $\beta(2) = 4$ which completes prof of conjectures by B. Duplantier et al [4].

References

[1] Applebaum D, *Levy Processes - From Probability to Finance and Quantum Groups*, Notices of the AMS, 51 (11): 13361347 (2004)

[2] D. Beliaev, S. Smirnov, *Harmonic measure and SLE*, Commun. Math. Phys. 290, 577595 (2009).

[3] John Cardy, *SLE for theoretical physicists*, Ann.Phys. 318 (2005) 81-118

[4] Bertrand Duplantier, Thi Phuong Chi Nguyen, Thi Thuy Nga Nguyen, Michel Zinsmeister, *Coefficient estimates for whole-plane SLE processes*, `http://hal.inria.fr/hal-00609774`, (2011).

[5] B.Duplantier, *Conformally invariant fractals and potential theory*, Phys.Rev.Lett., 84(7): 1363-1367 ,(2000).

[6] Ilya A. Gruzberg, *Stochastic geometry of critical curves, Schramm–Loewner evolutions and conformal field theory*, J. Phys. A: Math. Gen. **39**, no. 41 (2006) 12601–12655.

[7] T.C. Halsey, M.H. Jensen, L.P. Kadanoff, I. Procaccia, and B.I. Shraiman, *Fractal measures and their singularities: the characterization of strange sets*, Phys. Rev. A (3), 33(2): p. 1141, 1986.

[8] Matthew B. Hastings, *Exact Multifractal Spectra for Arbitrary Laplacian Random Walks*, *Phys. Rev. Lett.* **88** (2002) 055506.

[9] Gregory F. Lawler, *Conformally invariant processes in the plane*, Mathematical Surveys and Monographs, **114**, Amer. Math. Soc., Providence, RI, (2005).

[10] Gregory F. Lawler, *Conformal invariance and 2D statistical physics*, *Bull. Amer. Math. Soc.* **46** (2009) 35–54.

[11] Roelof Koekoek, Peter A. Lesky, René F. Swarttouw, *Hypergeometric orthogonal polynomials and their q-analogues*, Springer-Verlag, Berlin, 2010.

[12] Igor Loutsenko, *SLE$_\kappa$: correlation functions in the coefficient problem*, J. Phys. A: Math. Theor. 45 275001, 2012,(2012)

[13] Igor Loutsenko, Oksana Yermolayeva *Average Harmonic Spectrum of the Whole-Plane SLE*, BiBoS preprint 12-12-429 (2012), `http://www.math.uni-bielefeld.de/~bibos/preprints/12-12-429.pdf`

[14] Igor Loutsenko, Oksana Yermolayeva *On Exact Multifractal Spectrum of the Whole-Plane SLE*, `http://arxiv.org/abs/1203.2756` (2012)

[15] P. Oikonomou, I. Rushkin, I. A. Gruzberg, and L. P. Kadanoff, *Global properties of stochastic Loewner evolution driven by Lvy processes*, J. Stat. Mech. (2008) P01019

[16] I. Rushkin, P. Oikonomou, L. P. Kadanoff, and I. A. Gruzberg, *Stochastic Loewner evolution driven by Levy processes*, J. Stat. Mech. (2006) P01001